

Oral Biopsy Instructions

Specimen Handling



- Place specimen into appropriate jar containing 10% buffered formalin immediately.
- Specimens from more than 1 site must be placed in separate jars.
- Stick the patient label on the side of the jar (NOT on the lid).
- Indicate the specimen site on each container.

Requisition Completion

PMI Labs
Early diagnosis, better outcomes™

Perceptronix Medical Inc. - Vancouver
Suite 400, 555 West 8th Avenue, Vancouver, Canada V5Z 1C6
Suite 400, 555 West 8th Avenue, Vancouver, Canada V5Z 1C6
Hours of Operation: 8:00 am - 5:00 pm PST Tel: (604) 629-8785

HISTOLOGY REQUISITION FORM

Laboratory Use Only - Barcode Here

Patient Information	Physician Information
Medical # _____	Physician Name _____
Location _____	Specialty _____
Sex _____	DOB _____
Demographics _____	Phone _____
PHSA _____	Additional Physician _____
Specimen Site _____	Medical # _____
Specimen Container _____	
Formaldehyde _____	
Notes _____	
Clinical Summary	
Relevant Pathology _____	
Physician Signature _____	

- Place patient label on requisition (ensure jar label MATCHES the requisition label).
- Complete Physician Information section.
- Indicate number of sites such as A, B (ensure sites listed MATCH jars sent).
- Use the mouth diagram on the back to denote location and complete Clinical Summary section.

Specimen Return

- Place labelled jar into biohazard bag and completed requisition into bag pocket.
- Place sealed specimen bag into padded envelope or mailer box provided.
- Use padded envelope to send via Medi-Tran (MTS Logistics) or use return mailer box to send via Canada Post.